

Rocky Mountain Geology

Volume 35, Number 1, Spring, 2000

CONTENTS

	Page
<i>C. D. Frost</i>	1–5
<i>D. M. Cox, C. D. Frost, and K. R. Chamberlain</i>	7–30
<i>W. R. Premo and C. M. Fanning</i>	31–50
<i>W. R. Premo and R. R. Loucks</i>	51–70
<i>C. D. Frost, K. R. Chamberlain, B. R. Frost, and J. S. Scoates</i>	71–90
<i>C. S. Siddoway, R. M. Givot, C. D. Bodle, and M. T. Heizler</i>	91–111
<i>B. R. Edwards and C. D. Frost</i>	113–137

Reflections on Proterozoic magmatism of the Rocky Mountains and environs: Past and present
2.01-Ga Kennedy dike swarm, southeastern Wyoming: Record of a rifted margin along the southern Wyoming province
SHRIMP U-Pb zircon ages for Big Creek gneiss, Wyoming and Boulder Creek batholith, Colorado: Implications for timing of Paleoproterozoic accretion of the northern Colorado province
Age and Pb-Sr-Nd isotopic systematics of metaplutonic rocks from Green Mountain magmatic arc, southeastern Wyoming: Isotopic characterization of a Paleoproterozoic island arc system
The 1.76-Ga Horse Creek anorthosite complex, Wyoming: A massif anorthosite emplaced late in the Medicine Bow orogeny
Dynamic versus anorogenic setting for Mesoproterozoic plutonism in the Wet Mountains, Colorado: Does the interpretation depend on level of exposure?
An overview of the petrology and geochemistry of the Sherman batholith, southeastern Wyoming: Identifying multiple sources of Mesoproterozoic magmatism

UNIVERSITY OF WYOMING

ISSN 0010-7980